

12

EUROPEAN PATENT APPLICATION

21 Application number: 79301514.0

22 Date of filing: 30.07.79

51 Int. Cl.³: **C 07 H 17/08**
A 61 K 31/70
//A01N9/28

30 Priority: 02.08.78 US 930468

43 Date of publication of application:
 06.02.80 Bulletin 80/3

64 Designated Contracting States:
 AT BE CH DE FR GB IT LU NL SE

71 Applicant: **MERCK & CO. INC.**
 126, East Lincoln Avenue
 Rahway New Jersey 07065(US)

72 Inventor: **Fisher, Michael Herbert**
 1140 Concord Drive
 Bridgewater, New Jersey 08807(US)



72 Inventor: **Tolman, Richard Lee**
 29 Upper Warren Way
 Warren, New Jersey 07060(US)

74 Representative: **Crampton, Keith John Allen et al.**
 D YOUNG & CO 9 & 10 Staple Inn
 London WC1V 7RD(GB)

54 Sugar derivatives of C-076 compounds, their preparation and their use as antiparasitic agents.

57 The invention provides derivatives of C-076, which is a series of macrolides, in which the substituents are sugar or saccharide groups. The sugar groups are substituted on any of the available hydroxy groups of the C-076 molecule by various known procedures. The compounds thus produced, which have the structural formula:

in which the broken line indicates that a single or double bond is present in the 22(23) position; R₁ is hydroxy, loweralkanyloxy, or glycosyloxy, and is present only when the broken line indicates a single bond; R₂ is *iso*-propyl or *sec*-butyl; R₃ is hydrogen, methyl, loweralkanyloxy or glycosyl; and R is hydrogen, loweralkanyloxy, glycosyl,



where R₄ is hydrogen, loweralkanyloxy or glycosyl; in which the glycosyl groups are polyhydroxy (5 or 6)-membered cyclic acetals that are optionally substituted with lower alkyl and in which the hydroxy groups may be optionally substituted with lower alkyl or loweralkanyloxy and their mono-unsaturated derivatives; provided that at least one of R, R₁, R₂ or R₄ must contain a glycosyl group, and that if only R contains a glycosyl group, that group is other than α-L-oleandrosyl or α-L-oleandrosyl-α-L-oleandrosyl, have profound anthelmintic, insecticidal, ectoparasiticide and acaricidal activity and compositions for such uses are also disclosed.


EP 0 007 812 A1

SUGAR DERIVATIVES OF C-076 COMPOUNDS, THEIR PRE-
PARATION AND THEIR USE AS ANTIPARASTIC AGENTS


The term C-076 is used to describe a series of compounds isolated from the fermentation broth of a C-076-producing strain of Streptomyces avermitilis. The morphological characteristics of the culture are completely described in copending U.S. Patent Application Serial No. 772,601. The C-076 compounds are a series of structurally related macrolides, with one or more hydroxy substituents. The present invention is based on the discovery that the hydroxy substituents of C-076 and certain of its derivatives are capable of having, as substituent, a sugar molecule, and of procedures for the selective glycosylation at the various positions of such compounds having more than one hydroxy group.

Glycosylated compounds of the present invention have been found to possess profound anthelmintic, insecticidal, ectoparasitocidal and acaricidal activity.

The C-076 series of compounds have the following structural formula :



where R is the α -L-oleandrosyl- α -L-oleandrosyl group, which has the structural formula :



- 5 and where the broken line between C₂₂ and C₂₃ indicates that a single or a double bond is present in that position; R_1 is hydroxy

-3-

16024

and is present only when said broken line indicates a single bond;

R_2 is iso-propyl or sec-butyl; and

R_3 is methoxy or hydroxy.

5 There are eight different C-076 compounds and they are given the designation Ala, Alb, A2a, A2b, Bla, Blb, B2a, B2b based upon the structure of the individual compounds.

10 In the foregoing structural formula, the individual C-076 compounds are as set forth below.

	R_1	R_2	R_3
Ala	Double bond	<u>sec</u> -butyl	-OCH ₃
Alb	Double bond	<u>iso</u> -propyl	-OCH ₃
A2a	-OH	<u>sec</u> -butyl	-OCH ₃
15 A2b	-OH	<u>iso</u> -propyl	-OCH ₃
Bla	Double bond	<u>sec</u> -butyl	-OH
Blb	Double bond	<u>iso</u> -propyl	-OH
B2a	-OH	<u>sec</u> -butyl	-OH
B2b	-OH	<u>iso</u> -propyl	-OH

20 As is readily seen, all of the C-076 compounds have hydroxy groups at the 7-position and the 4"-position of the carbohydrate side chain. Thus all of the compounds have at least two hydroxy groups capable of being glycosylated. In addition, 25 the A2 and B1 series of compounds have a third hydroxy group and the B2 series of compounds has a fourth hydroxy group which may be glycosylated.

The carbohydrate side chain may also be hydrolyzed to remove one or both of the α -L-oleandrose groups. In this case there would remain a hydroxy 30 group at the 4' or 13-position with the removal of a single α -L-oleandrose (monosaccharide) or both α -L-oleandrose (aglycone) respectively. These hydroxy groups may readily be glycosylated.

-4-

16024

The monosaccharide and aglycone derivatives are prepared by the treatment of the parent C-076 compound with acid. The outer α -L-oleandrose group is more easily removed than the α -L-oleandrose group directly bonded to the C-076 substrate thus facilitating the separate preparation of the monosaccharide and aglycone without contamination with the other reaction product.


The process employed for the removal of the C-076 carbohydrate group or groups is to put the C-076 starting material in solution in a mixture of from 0.01 to 10% acid in a non-nucleophilic water miscible solvent such as dioxane, tetrahydrofuran, dimethoxyethane, dimethylformamide, bis-2-methoxyethyl ether and the like, and from 0.1 to 20% water. The mixture is stirred for from 6 to 24 hours at room temperature to complete the reaction. Acids such as sulfuric, hydrochloric, hydrobromic, phosphoric, trifluoroacetic and trifluorosulfonic are acceptable. Sulfuric acid is preferred.

When lower acid concentrations such as from 0.01 to 0.1% are employed, the monosaccharide is predominantly prepared. When higher concentrations of acid are employed, such as in the range of 1 to 10%, the aglycone is predominantly prepared. Intermediate concentrations of acid will tend to prepare mixtures of monosaccharide and aglycone which are generally separable using chromatographic techniques.

The monosaccharide may also be prepared by stirring the C-076 precursor for from 6 to 24 hours at room temperature in 1% sulfuric acid in isopropanol. In addition the aglycone can be prepared by stirring the C-076 precursor for from 6 to 24 hours at room temperature in 1% sulfuric acid in methanol. The other acids listed above may also be employed in this

process. This process is preferred for use with the 2- series of C-076 compounds since some addition may be observed to the 22,23 double bond of the series of C-076 compounds with a 22,23 unsaturation. The desired monosaccharide or aglycone are
5 isolated and purified using known techniques.

The glycosyl compounds of this invention have the following structural formula :



where the broken line indicates that a single or a double bond
10 is present in the 22(23) position;


R₁ is hydroxy, loweralkanoyloxy or glycosyloxy and is present only when the broken line indicates a single bond;

R₂ is iso-propyl or sec-butyl;

R₃ is hydrogen, methyl, loweralkanoyl or glycosyl;

15 and

R is hydrogen, loweralkanoyl, glycosyl,



where R_4 is hydrogen, loweralkanoyl or glycosyl; provided that at least one of R , R_1 , R_3 or R_4 must contain a glycosyl group, and that if only R contains a glycosyl group, that glycosyl group is other than α -L-oleandrosyl or α -L-oleandrosyl- α -L-oleandrosyl. The glycosyl groups are polyhydroxy (5 or 6)-membered cyclic acetals that are optionally substituted with lower alkyl and in which the hydroxy groups may be optionally substituted with lower alkyl or lower alkanoyl and their mono-unsaturated derivatives.

In the present specification the term "loweralkanoyl" means alkanoyl groups of from 2 to 10 carbon atoms such as acetyl, propionyl, butyryl, pentanoyl, hexanoyl, pivaloyl, octanoyl and decanoyl. "Lower alkyl" means " C_{1-10} alkyl", but preferably C_{1-6} alkyl such as methyl, ethyl, propyl, isopropyl or a butyl.

The nature of the sugar residue in the above glycosyl groups is not critical and any sugar may be substituted onto the C-076 substrate using the procedures described below. The preferred sugar residues are glucopyranosyl, galactopyranosyl, mannopyranosyl, maltosyl, arabinopyranosyl, lyxopyranosyl, xylopyranosyl, ribopyranosyl, oleandrosyl, rhamnopyranosyl, fucopyranosyl, lactosyl, ribofuranosyl, mannofuranosyl, glucofuranosyl, arabinofuranosyl, mycarosyl, cladinosyl, desosaminosyl, daunosaminosyl, mycaminosyl, cymarosyl and olivosyl.

The foregoing sugars are available generally in the D or L configuration. The instant invention includes both the possible configurations for attachment to the C-076 substrate.

5 The above carbohydrate or sugar groups may be substituted on the C-076 compounds as mono, di or tri- saccharides wherein one of the above sugar groups is further substituted with another of the same or different sugar. In addition, where there
10 is more than one hydroxy group available for substitution, the sugar groups may be present on only one or on more than one of such hydroxy groups, and the substitution may be with identical or different sugar moieties.

15 The preferred sugar substituents are mono- or disaccharide substitution with glucopyranosyl, rhamnopyranosyl, oleandrosyl or olivosyl groups. The most preferred groups are glucopyranosyl and oleandrosyl.

20 The processes for the substitution of the carbohydrate groups onto the hydroxy groups of the substrate molecule the Koenigs-Knorr process, the silver triflate process the orthoester process, or the glycol addition process.

25 The carbohydrate starting materials employed for the Koenigs-Knorr, the Helferich modification thereof and the silver triflate processes are protected by acylating all of the free hydroxy groups. The preferred protecting group is the acetyl
30 group, however, other groups such as the benzoate may be employed. The processes for the blocking of the hydroxy groups are well known to those skilled in the art. The acetyl blocking groups are also easily removed at the completion of the reaction by
35 catalytic hydrolysis, preferably base-catalyzed hydrolysis such as with an alcoholic ammonia solution.

The Koenigs-Knorr and silver triflate processes use as starting materials the acetohalosugars preferably the acetobromo sugars such as the appropriate acetobromohexoses and acetobromopentoses of the sugar groups listed above. The bromine atom is substituted on a carbon atom adjacent to an acetyl group and the sugar moiety become bonded to the substrate at the carbon atom to which the halogen was attached.

10 In the Koenigs-Knorr reaction the C-076 compound is dissolved under anhydrous conditions in an aprotic solvent. Ether is the preferred solvent, however, methylene chloride, acetonitrile, nitromethane, dimethoxy ethane and the like may also
15 be employed. To the substrate solution is added the acetohalosugar and silver oxide. A single molar equivalent of the sugar is required for the reaction, however, an additional 10 to 15 moles occasionally
20 aids the reactions. Additional molar excesses beyond 15 may be employed in difficult reactions, however, such very large excesses tend to make the isolation of the product more difficult. It has been found preferable to employ freshly prepared silver oxide for the reaction, since the material
25 tends to lose some of its catalytic efficiency upon standing for prolonged periods. The silver oxide is prepared from silver nitrate using known procedures. The reaction may be carried out at from 10-15°C, however, reaction at room temperature is
30 preferred. The reaction generally requires from 2 to 10 days for completion. Reaction progress is monitored by taking aliquots from the reaction mixture and examining them with thin layer chromatographic techniques. Possible side reactions are
35 avoided by carrying out the reaction in the dark,

and this method is preferred. The product is isolated using techniques known to those skilled in the art.

In one modification of the Koenigs-Knorr reaction, known as the Helferich modification thereof, a mercuric halide, such as mercuric chloride or bromide, alone or in combination with mercuric oxide or mercuric cyanide is substituted for the silvoxide. The above described reaction conditions be employed except that nitromethane and benzene are the preferred solvents and reflux temperature is the preferred reaction temperature.

The silver triflate reaction uses the reagent silver triflate (silver trifluoromethyl sulfonate) and the acetohalosugar in the same solvents listed above, with ether being preferred. The silver triflate is best if highly purified and freshly prepared just prior to its use. Methods for the preparation of silver triflate are well known to those skilled in the art. All of the reactants are combined in the solvent and the reaction conducted at from 10 to 50°C for from 2 to 48 hours. Generally, however, the reaction is complete in about 24 hours at room temperature. The progress of the reaction may be followed by thin layer chromatography techniques. Again the reaction is preferably carried out in the dark, and with absolutely dry reactants and equipment.

A single mole of the sugar is required, however, a single molar excess is often used to aid in the course of the reaction.

During the course of the reaction a mole of triflic acid (trifluoromethanesulfonic acid) is liberated. This is a very strong acid and a molar

equivalent of a base is required to neutralize the acid. Preferred bases are non-nucleophilic bases such as tertiary amines, preferably triethylamine, diisopropylethylamine, diazabicyclo-undecane, diazabicyclononane and the like. Since triflic acid is such a strong acid, if the base used is not a strong enough base to neutralize all of the acid, the residual acid will adversely affect the course of the reaction and of the isolation of the product. The product is isolated using techniques known to those skilled in the art.

The orthoester process prepares sugar derivatives of the C-076 compounds from ortho-esters of a lower alkanol and of the above sugars at the hydroxy function of said C-076 compounds. The ortho-esters are prepared from the acetohalosugars using a lower alkanol and procedures which are well known to those skilled in the art. The reaction is carried out in an aprotic solvent such as dichloroethane, nitromethane, methylene chloride, dimethoxy ether, acetonitrile, tetrahydrofuran and the like. Dichloroethane, nitromethane, dimethoxy ethane and tetrahydrofuran are preferred. The reaction is preferably carried out at the reflux temperature of the reaction mixture and is generally complete in from about 4 to 24 hours. Catalytic amounts of mercuric bromide or mercuric chloride are added to aid in the reaction. During the course of the reaction one mole of the alcohol used to make the orthoester is liberated. Thus, the preferred method is to azeotropically distill off the solvent to remove the alcohol and to force the reaction to completion. To prevent any volume reduction, fresh solvent is added as the distillation proceeds to maintain a constant volume. To isolate

the product, the solvent is generally removed and the residue washed with a reagent to remove the mercury salts, such as aqueous potassium iodide. The product is then isolated using known techniques.

- 5 Another glycosylation procedure has been developed which employs glycal starting materials. Glycals are 1,2-unsaturated cyclic sugars where the ring is a 6 membered ring. The reaction provides for the substitution of the glycosyl moiety onto the
- 10 C-076 substrate in two ways, depending on the substitution at the 3-position of the sugar ring. Where the 3-position is substituted with a leaving group, such as a loweralkanoyloxy, such as acetate, substituted phenylsulfonate, such as a tosylate,
- 15 the leaving group is eliminated and the double bond shifts to the 2,3-position and the 3-position in the resultant product is unsubstituted.

- If the 3-position group is not a leaving group, such as loweralkoxy, preferably methoxy, the
- 20 group is not eliminated and the 3-position remains intact. The reaction produces an addition across the double bond, and the resultant saturated product is attached to the C-076 substrate at the 1-position.

- The reaction is carried out in an aprotic,
- 25 non-polar solvent or, since the glycal starting materials are often liquids at room temperature, the reaction may be carried out neat. Preferred solvents are halogenated hydrocarbons such as methylene chloride, chloroform and the like, acetonitrile,
- 30 benzene, toluene, ether and the like. In addition, the reaction may contain a catalytic amount of a Lewis Acid such as boron trifluoride, tin chloride (SnCl_2) titanium, tetrachloride, zinc chloride (ZnCl_2) and the like or other acid catalyst (β -
- 35 toluenesulfonic acid, pyridinium β -toluenesulfonate).

-12-

16024

The reaction is carried out at from room temperature to the reflux temperature of the reaction mixture, however usually maximum temperatures of 75°C are adequate. The reaction is
5 complete in from 1/2 to 24 hours and the product is isolated using known techniques.

With respect to the foregoing glycosylation reactions, it is noted that the B-series of compounds are preferably glycosylated using the Koenigs-Knorr
10 or glycal addition reaction. This is because the B-compounds are somewhat sensitive to the mercury compounds used in the Helferich modification.

Because of the different reactivities of the various hydroxy groups, and with the use of certain
15 protecting techniques, all possible combinations of glycosylated C-076 compounds can be prepared. The hydroxy groups at the 5 and 23 positions are much more reactive to glycosylation than the 4", 4' or
20 13 positions. Thus, where the 5 or 23 glycosylated products are desired, no protection is necessary for the 4", 4' or 13 positions.

The C-076 A1 compounds have only the 4", 4' or 13 positions available for glycosylation, thus no reaction is possible at the 5 or 23 positions.

25 The C-076 A2 compounds have only the 23-hydroxy group available for glycosylation, in addition to the 4", 4' or 13 positions. Thus, the 23 glycosyl compound can be prepared with no protection reactions necessary. The 4", 4' or 13
30 positions can be glycosylated by protecting the 23-position following the procedure described below.

The C-076 B1 compounds have only the 5-position available in addition to the 4", 4' or 13 positions. If the 5-glycosyl compounds are
35 desired, the reaction is carried out without the protection of the 4", 4' or 13 positions. If the

-13-

16024

4", 4' or 13 glycosyl compounds are desired, protection of the 5-position following the procedure described below is required.

The C-076 B2 compounds have both the 5 and 23 positions available for glycosylation in addition to the 4", 4' or 13 positions. The reactivity of the 5- and 23-position hydroxy groups are about equal. If glycosylation at the 5 and 23 positions is desired, the reaction is carried out as above described. If glycosylation at the 4", 4' or 13 positions is desired, protection of both of the 5 and 23 hydroxy groups is necessary. If glycosylation of only one of the 5- or 23-positions is desired, the reaction is carried out as above described using minimal times and temperatures within the ranges given, thus resulting in a mixture of 5- and 23-glycosyl compounds. Such a mixture is readily separated, usually with chromatographic techniques such as column, high pressure, liquid chromatography and thin layer or preparative layer chromatography. The technique of thin layer chromatography is usually employed to follow the course of the reaction in order to maximize the formation of the individual 5- and 23-substituted compounds and to avoid the formation of the 5,23-disubstituted compounds.

The protection of the 5 and 23 positions is carried out in two ways; one applicable to the 5 and 23 positions, and the other applicable to the 5-position only.

The 5 and/or the 23 positions may be readily protected by acylating, preferably acetylating. The C-076 compound to be protected is dissolved in an aprotic, non polar solvent, such as one of such solvents described above preferably

-14-

16024

ether and an acyl halide, preferably acetyl chloride, is added dropwise substantially at room temperature but up to a 10-40°C range. The reaction mixture is stirred for from 2 to 6 hours. A catalytic amount of silver oxide (Ag_2O) is added to the reaction. The acylated compound is isolated using known techniques.

The acyl protecting group may be removed by base catalyzed hydrolysis, such as with an alkali metal alkoxide, preferably sodium methoxide, in the corresponding alcohol, preferably methanol.

In addition, the 5-position alcohol may be oxidized to the ketone using manganese dioxide (MnO_2) in ether. The reaction is completed in about 10 to 30 hours at about room temperature.

The ketone is then readily converted back to the hydroxy group using borohydride reduction, preferably sodium borohydride. The reaction is complete in about 5 minutes to 1 hour with stirring substantially at room temperature.

This oxidation-reduction reaction sequence is seen to aid in the preparation of C-076 B2 compounds with only one of the 5- or 23-positions glycosylated. By protecting the 5-position with the preparation of the 5-ketone the 23-glycosyl compound may readily be prepared. Or the 23-acyl-5-ketone compound can be prepared, the 5-position reduced to the hydroxy group, and the 5-glycosyl compound prepared.

It is noted that the acyl derivatives are included within the ambit of this invention, thus, if desired the acyl protecting groups need not be removed at the end of a reaction sequence. In addition, once the glycosyl compounds are prepared, at the desired position or positions,

-15-

16024

any remaining hydroxy groups may be acylated. The foregoing procedure may be employed to acylate the 5- and 23-positions. Where glycosylation has occurred or is planned for the 5- and/or 23-positions the 4", 4' or 13 position may be acylated by dissolving the compound in a suitable solvent, preferably pyridine, and adding the acylating reagent, preferably a loweralkanoyl halide, such as acetyl chloride, dropwise. The reaction is maintained at 10 from 0°C to room temperature for from 4 to 24 hours. The product is isolated using known techniques.

The novel glycosylated compounds of this invention have significant parasitocidal activity as anthelmintics, insecticides, ectoparasitocides 15 and acaricides, in human and animal health and in agriculture.

The disease or group of diseases described generally as helminthiasis is due to infection of an animal host with parasitic worms known as helminths. 20 Helminthiasis is a prevalent and serious economic problem in domesticated animals such as swine, sheep, horses, cattle, goats, dogs, cats and poultry. Among the helminths, the group of worms described as nematodes causes widespread and often times serious 25 infection in various species of animals. The most common genera of nematodes infecting the animals referred to above are Haemonchus, Trichostrongylus, Ostertagia, Nematodirus, Cooperia, Ascaris, Bunostomum, Oesophagostomum, Chabertia Trichuris, 30 Strongylus, Trichonema, Dictyocaulus, Capillaria, Heterakis, Toxocara, Ascaridia, Oxyuris, Ancylostoma, Uncinaria, Toxascaris and Parascaris. Certain of these, such as Nematodirus, Cooperia, and Oesophagostomum attack primarily the intestinal 35 tract while others, such as Haemonchus and

-16-

16024

Ostertagia, are more prevalent in the stomach while still others such as Dictyocaulus are found in the lungs. Still other parasites may be located in other tissues and organs of the body such as the heart and blood vessels, subcutaneous and lymphatic tissue and the like. The parasitic infections known as helminthiases lead to anemia, malnutrition, weakness, weight loss, severe damage to the walls of the intestinal tract and other tissues and organs and, if left untreated, may result in death of the infected host. The glycosylated C-076 compounds of this invention have unexpectedly high activity against these parasites, and in addition are also active against Dirofilaria in dogs, Nematospirides, Syphacia, Aspiculuris in rodents, arthropod ectoparasites of animals and birds such as ticks, mites, lice, fleas, blowfly, in sheep Lucilia sp., biting insects and such migrating dipterous larvae as Hypoderma sp. in cattle, Gastrophilus in horses, and Cuterebra sp. in rodents.

The instant compounds are also useful against parasites which infect humans. The most common genera of parasites of the gastro-intestinal tract of man are Ancylostoma, Necator, Ascaris, Strongyloides, Trichinella, Capillaria, Trichuris, and Enterobius. Other medically important genera of parasites which are found in the blood or other tissues and organs outside the gastrointestinal tract are the filarial worms such as Wuchereria, Brugia, Onchocerca and Loa, Dracunculus and extra intestinal stages of the intestinal worms Strongyloides and Trichinella. The compounds are also of value against arthropods parasitizing man, biting insects and other dipterous pests causing annoyance to man.

-17-

16024

The compounds are also active against household pests such as the cockroach, Blatella sp., clothes moth, Tineola sp., carpet beetle, Attagenus sp. and the housefly Musca domestica.

- 5 The compounds are also useful against insect pests of stored grains such as Tribolium sp., Tenebrio sp. and of agricultural plants such as spider mites, (Tetranychus sp.), aphids, (Acyrtiosiphon sp.); against migratory orthopterans such as locusts and
- 10 immature stages of insects living on plant tissue. The compounds are useful as a nematocide for the control of soil nematodes and plant parasites such as Meloidogyne spp. which may be of importance in agriculture.
- 15 These compounds may be administered orally in a unit dosage form such as a capsule, bolus or tablet, or as a liquid drench where used as an anthelmintic in mammals. The drench is normally a solution, suspension or dispersion of the active
- 20 ingredient usually in water together with a suspending agent such as bentonite and a wetting agent or like excipient. Generally, the drenches also contain an antifoaming agent. Drench formulations generally contains from about 0.001 to 0.5% by weight
- 25 of the active compound. Preferred drench formulations may contain from 0.01 to 0.1% by weight. The capsules and boluses comprise the active ingredient admixed with a carrier vehicle such as starch, talc, magnesium stearate, or di-calcium phosphate.
- 30 Where it is desired to administer the C-076 compounds in a dry, solid unit dosage form, capsules, boluses or tablets containing the desired amount of active compound usually are employed. These dosage forms are prepared by intimately and uniformly mixing
- 35 the active ingredient with suitable finely divided

-18-

16024

diluents, fillers, disintegrating agents and/or binders such as starch, lactose, talc, magnesium stearate, vegetable gums and the like. Such unit dosage formulations may be varied widely with respect to their total weight and Content of the antiparasitic agent depending upon factors such as the type of host animal to be treated, the severity and type of infection and the weight of the host.

When the active compound is to be administered via an animal feedstuff, it is intimately dispersed in the feed or used as a top dressing or in the form of pellets which may then be added to the finished feed or optionally fed separately. Alternatively, the antiparasitic compounds of our invention may be administered to animals parenterally, for example, by intraruminal, intramuscular, intratracheal, or subcutaneous injection in which event the active ingredient is dissolved or dispersed in a liquid carrier vehicle. For parenteral administration, the active material is suitably admixed with an acceptable vehicle, preferably of the vegetable oil variety such as peanut oil, cotton seed oil and the like. Other parenteral vehicles such as organic preparations using solketal, glycerol-formal and aqueous parenteral formulations are also used. The active acylated C-076 compound or compounds are dissolved or suspended in the parenteral formulation for administration; such formulations generally contain from 0.005 to 5% by weight of the active compound.

Although the antiparasitic agents of this invention find their primary use in the treatment and/or prevention of helminthiasis, they are also useful in the prevention and treatment of diseases

caused by other parasites, for example, arthropod parasites such as ticks, lice, fleas, mites and other biting and sucking insects in domesticated animals and poultry. They are also effective

5 in treatment of parasitic diseases that occur in other animals including humans. The optimum amount to be employed for best results will, of course, depend upon the particular compound employed, the species of animal to be treated and the type and severity of

10 parasitic infection or infestation. Generally, good results are obtained with our novel compounds by the oral administration of from about 0.001 to 10 mg. per kg. of animal body weight, such total dose being given at one time or in divided doses

15 over a relatively short period of time such as 1-5 days. With the preferred compounds of the invention, excellent control of such parasites is obtained in animals by administering from about 0.025 to 0.5 mg. per kg. of body weight in a single dose. Repeat

20 treatments are given as required to combat re-infections and are dependent upon the species of parasite and the husbandry techniques being employed. The techniques for administering these materials to animals are known to those skilled in the

25 veterinary field. The compounds may also be administered in combination with other antiparasitic compounds or compounds with other biological activities to provide for a single treatment with a broadened spectrum of activity.

30 When the compounds described herein are administered as a component of the feed of the animals, or dissolved or suspended in the drinking water, compositions are provided in which the active compound or compounds are intimately dispersed in an inert

35 carrier or diluent. By inert carrier is meant one

that will not react with the antiparasitic agent and one that may be administered safely to animals. Preferably, a carrier for feed administration is one that is, or may be, an ingredient of the animal
5 ration.

Suitable compositions include feed premixes or supplements in which the active ingredient is present in relatively large amounts and which are suitable for direct feeding to the animal or for
10 addition to the feed either directly or after an intermediate dilution or blending step. Typical carriers or diluents suitable for such compositions include, for example, distillers' dried grains, corn meal, citrus meal, fermentation residues, ground
15 oyster shells, wheat shorts, molasses solubles, corn cob meal, edible bean mill feed, soya grits, crushed limestone and the like. The active glycosylated C-076 compounds are intimately dispersed throughout the carrier by methods such as grinding, stirring,
20 milling or tumbling. Compositions containing from about 0.005 to 2.0% by weight of the active compound are particularly suitable as feed premixes. Feed supplements, which are fed directly to the animal, contain from about 0.0002 to 0.3% by weight
25 of the active compounds.

Such supplements are added to the animal feed in an amount to give the finished feed the concentration of active compound desired for the treatment and control of parasitic diseases. Although
30 the desired concentration of active compound will vary depending upon the factors previously mentioned as well as upon the particular glycosylated C-076 compound employed, the compounds of this invention are usually fed at concentrations of between 0.00001 to

-21-

16024

0.002% in the feed in order to achieve the desired antiparasitic result.

In using the compounds of this invention, the individual glycosylated C-076 components may be prepared and used in that form. Alternatively, mixtures of two or more of the individual glycosylated C-076 compounds may be used.

In the isolation of the C-076 compounds, which serve as starting materials for the instant processes, from the fermentation broth, the various C-076 compounds will be found to have been prepared in unequal amounts. In particular an "a" series compound will be prepared in a higher proportion than the corresponding "b" series compound. The weight ratio of "a" series to the corresponding "b" series is about 85:15 to 99:1. The differences between the "a" series and "b" series is constant throughout the C-076 compounds and consists of a butyl group and a propyl group respectively at the 25 position. This difference, of course, does not interfere with any of the instant reactions. In particular, it may not be necessary to separate the "b" components from the related "a" component. Separation of these closely related compounds is generally not practiced since the "b" compound is present only in a very small percent by weight, and the structural difference has negligible effects on the reaction processes and biological activities.

The C-076 compounds of this invention are also useful in combatting agricultural pests that inflict damage upon crops while they are growing or while in storage. The compounds are applied using known techniques as sprays, dusts, emulsions and the like, to the growing or stored crops to effect

-22-

16024

protection from such agricultural pests.

The following examples are provided in order that the invention might be more fully understood; they are not to be construed as
5 limitations of the invention.

The C-076 glycosyl derivatives prepared in the following examples are generally isolated as amorphous solids and not as crystalline solids. They are thus characterized analytically using
10 techniques such as mass spectrometry, nuclear magnetic resonances and the like. Being amorphous, the compounds are not characterized by sharp melting points, however the chromatographic and analytical methods employed indicate that the compounds are
15 pure.

EXAMPLE 1

4'-O-(2,3,4,6-tetra-O-acetyl-β-D-glucopyranosyl)
C-076 A_{2a} monosaccharide

A 3-necked flask is charged with C-076
20 A_{2a} monosaccharide (50 mg.), 3,4,6-tri-O-acetyl-
1,2-O-(ethylorthoacetyl)-X-D-glucopyranose (50 mg.),
mercuric bromide (2 mg.) and dichloroethane (20 ml.).
The flask is fitted into a dropping funnel containing
fresh, dry solvent and a Dean-Stark Trap. The
25 mixture is heated at reflux under nitrogen and as
the solvent is slowly distilled from the reaction
mixture, it is replaced with fresh solvent from the
dropping funnel. After 24 hours of refluxing, thin
layer chromatographic monitoring indicates no further
30 reaction progress and the mixture is cooled,
filtered (discarding all solids), and the filtrate
evaporated to dryness in vacuo. Chromatographic

-23-

16024

resolution of the mixture on preparative layer chromatography [silica gel, multiple development with benzene/2-propanol (19:1)] affords a large amount of starting material, some minor side products and 16 mg. of 4'-O-(2,3,4,6-tetra-O-acetyl- β -D-glucopyranosyl) C-076 A_{2a} monosaccharide, which is lyophilized from benzene. The structure is consistent with mass spectral and 300 MHz nuclear magnetic resonance analysis.

10

EXAMPLE 2

4''-O-(2,3,4,6-tetra-O-acetyl- β -D-glucopyranosyl)
C-076 B_{1a}

A 3-necked flask is charged with C-076 B_{1a} (50 mg.), 3,4,6-tri-O-acetyl-1,2-O-(ethyl-orthoacetyl)- α -D-glucopyranose (35 mg.), mercuric bromide (2 mg.) and dichloroethane (20 ml.). The reaction mixture is refluxed following the procedure of Example 1. After 12 hours of refluxing, a product forms which is isolated by cooling the reaction mixture, filtering and discarding all solids and evaporating the filtrate in vacuo. Chromatographic resolution of the mixture on preparative layer chromatography [silica gel, multiple development with 19:1 (benzene 12-propanol)] affords a large amount of starting material, small amounts of presumably isomeric glycosides and 3 mg. of 4''-O-(2,3,4,6-tetra-O-acetyl- β -D-glucopyranosyl) C-076 B_{1a}. The structure proposed is consistent with mass spectral and 300 MHz nuclear magnetic resonance analysis.

EXAMPLE 3

23-O-(2,2',3,3',4',6,6'-hepta-O-acetyl-β-maltosyl)

C-076 A_{2a}

To C-076 A_{2a} (20 mg.) in anhydrous ether
 5 (50 ml.) is added freshly prepared silver oxide
 (60 mg.) and 2,2',3,3',4',6,6'-hepta-O-acetyl-α-
 acetyl-α-maltosyl bromide (51 mg.). The sealed
 flask is stirred magnetically in the dark. After
 24 hours, another 20 mg. of 2,2',3,3',4',6,6'-
 10 hepta-O-acetyl-β-maltosyl bromide is added. After
 a total of five days, the reaction progress is
 judged to have stopped, the solids are filtered,
 washed with ether and discarded. The filtrate is
 evaporated under a stream of nitrogen and fractionated
 15 using preparative layer chromatography on silica
 gel using multiple development with 2-propanol
 /benzene (1:19). In addition, to some starting
 material and other by-products there is obtained
 21 mg. of 23-O-(2,2',3,3',4',6,6'-hepta-O-acetyl-β-
 20 maltosyl C-076 A_{2a} (lyophilized from benzene) which
 is judged to be homogeneous by nuclear magnetic
 resonance and thin layer chromatography. Mass
 spectral and 300 MHz nuclear magnetic resonance
 data are consistent with the product structure.

25

EXAMPLE 4

13-O-(4-O-acetyl-α-L-erythro-2-hexenopyranosyl)

C-076 A_{2a} Aglycon

C-076 A_{2a} aglycon (50 mg.) and 3,4-di-O-
 acetyl-L-rhamnal (50 mg.) are dissolved in benzene
 30 (20 ml.), a trace of p-toluenesulfonic acid is added
 and the mixture is stirred at 80°C. After 4 days
 the reaction is complete (no more starting material

-25-

16024

is present by thin layer chromatography). The solvent is evaporated in vacuo and chromatographic resolution of the products [preparative layer chromatography, silica gel using benzene 2-propanol (19:1) multiple development affords one major band which is subsequently resolved into 2 components (using the same separation procedure). By mass spectral and 300 MHz nuclear magnetic resonance analysis the two products are identified as 13-Q-
5 (4-Q-acetyl- α -L-erythro-2-hexenopyranosyl) C-076
10 A_{2a} aglycon (19 mg.) and 13,23-bis-(4-Q-acetyl- α -L-erythro-2-hexenopyranosyl) C-076 A_{2a} aglycon (11 mg.).

EXAMPLE 5

23-Q-acetyl-4"-Q-(2,3,4,6-tetra-Q-acetyl- β -D-gluco-
15 pyranosyl) C-076 A_{2a} .

To 23-Q-acetyl-C-076 A_{2a} (20 mg.) in anhydrous ether (20 ml.) is added 2,3,4,6-tetra-Q-acetyl- α -D-glucopyranosyl bromide (70 mg.) and freshly prepared silver oxide (200 mg.). The
20 sealed flask is stirred magnetically in the dark. After four days thin layer chromatography monitoring fails to show any further change in the reaction mixture composition and the solids are filtered, washed with ether and discarded.
25 The filtrate is evaporated under a stream of nitrogen and fractionated with preparative layer chromatography (silica gel) with multiple development using 19:1 (benzene/2-propanol).
23-Q-acetyl-4"-Q-(2,3,4,6-tetra-Q-acetyl- β -D-gluco-
30 pyranosyl) C-076 A_{2a} (6 mg.) is isolated after

-26-

16024

lyophilization from benzene in addition to other minor products and a considerable amount of starting material. Mass spectral and 300 MHz nuclear magnetic resonance analysis of the product are consistent with the structure.

EXAMPLE 6

5-Q-acetyl-4"-Q-(2,3,4,6-tetra-Q-acetyl- β -D-glucopyranosyl) C-076 B_{1a}

To C-076 B_{1a} (20 mg.) in anhydrous ether (8 ml.) is added freshly prepared silver oxide (23 mg.) and acetyl chloride (9.5 mg.). The reaction vessel is stirred magnetically under nitrogen in the dark until monoacetylation (at the 5-position) is complete (3-4 hours). Solid powdered sodium bicarbonate is added and the mixture is stirred for an additional hour. The solids are centrifuged and the supernatant containing 5-Q-acetyl C-076 B_{1a} with traces of C-076 B_{1a} and 5,4"-di-Q-acetyl C-076 B_{1a} (by thin layer chromatography) is poured onto 2,3,4,6-tetra-Q-acetyl- α -D-glucopyranosyl bromide (25 mg.) and freshly prepared silver oxide (25 mg.). After several days no further reaction progress is noted by thin layer chromatography [two subsequent additions of halogenose and silver oxide (25 mg. each) are made] after 24 and 48 hours of reaction. The solids are filtered, washed with ether and discarded. The filtrate evaporated under a stream of nitrogen and resolved into macrolide components and sugar decomposition products on a short overloaded

-27-

16024

silica gel column eluting with dichloromethane/
methanol (19:1). The macrolide components are
resolved on silica gel preparative layer
chromatography using multiple development with
5 benzene/isopropanol (95:5). The product is
isolated by lyophilization from benzene. Mass
spectral and 300 MHz nuclear magnetic resonance
data are consistent with the structure of
5-Q-acetyl-4"-Q-(2,3,4,6-tetra-Q-acetyl- β -D-gluco-
10 pyranosyl) C-076 B_{1a}.

EXAMPLE 7

13-Q-(2,3,4,6-tetra-Q-acetyl-D-glucopyranosyl)
A_{2a} aglycone

A 3-necked flask is charged with C-076
15 A_{2a} aglycone (30 mg.), 3,4,6-tri-Q-acetyl-1,2-Q-
(ethylorthoacetyl)- α -D-glucopyranose (50 mg.),
dichloroethane (50 ml.) and mercuric bromide
(1 mg.). The flask is fitted with a dropping
funnel containing fresh solvent and a Dean-Stark
20 Trap. The mixture is heated at reflux under
nitrogen as in Example 1. After 40 hours of
refluxing, monitoring by thin layer chromatography
indicates no further reaction progress and the
mixture is evaporated in vacuo. Resolution on
25 preparative layer chromatography [silica gel,
multiple development with benzene (2-propanol (19:1))]
affords a large amount of starting material (31 mg.)
and a slower moving band (6 mg.) which is shown
by mass spectrometry and 300 MHz nuclear magnetic
30 resonance to be 13-Q-(2,3,4,6-tetra-Q-acetyl-D-gluco-
pyranosyl) A_{2a} aglycone.

EXAMPLE 85-Q-(2,3,4,6-tetra-Q-acetyl-D-galactopyranosyl)-C-076 B_{1a}

- C-076 B_{1a} (250 mg.), freshly prepared
5 silver oxide (750 mg.) and 2,3,4,6-tetra-Q-
acetyl- α -D-galactopyranosyl bromide (150 mg.) are
added to dry ether (100 ml.) in a magnetically-
stirred round bottom flask and stirred under
nitrogen in the dark for four days. Every 24 hours
10 another 150 mg. of silver oxide and 100 mg. of
2,3,4,6-tetra-Q-acetyl-D-galactopyranosyl-C-076
B_{1a} are added (total: 1150 mg. of silver oxide and
350 mg. of 2,3,4,6-tetra-Q-acetyl-D-galactopyranosyl
C-076 B_{1a}). The solids are then filtered and washed
15 with ether. The filtrate is evaporated in vacuo
and the residual gum is chromatographed on silica
gel (200 gm.) using dichloromethane/methanol
(99:1) as eluant. Starting material (C-076 B_{1a},
170 mg.) is recovered as well as an additional product-
20 containing band which is further purified by
preparative layer chromatography [multiple
development with benzene/2-propanol (19:1)] to
furnish the desired product. The product is
lyophilized from benzene; spectral analysis is
25 consistent with 5-Q-(2,3,4,6-tetra-Q-acetyl-D-
galactopyranosyl)-C-076 B_{1a}.

EXAMPLE 94'-Q-(3,4-di-Q-acetyl-2,6-di-deoxy-L-arabino-hexopyranosyl)-C-076 A_{2a} Monosaccharide

- A 110 ml. round-bottomed flask is charged with C-076 A_{2a} monosaccharide (40 mg.), 3,4-di-Q-acetyl-2,6-dideoxy- α -L-arabino-hexopyranosyl chloride (diacetyl-oliviosyl chloride, 91 mg.), mercuric bromide (216 mg.), mercuric cyanide (250 mg.), and dry ether (50 ml). The stoppered flask is stirred in the dark under nitrogen until the reaction appears to make no further progress (by thin layer chromatography; 6 days). Additional quantities of the 3,4-di-Q-acetyl-2,6-di-deoxy-L-arabino-hexopyranosyl)-chloride (2 X 63 mg.) are added after 24 and 48 hours. There are seven product bands in addition to a small amount of recovered starting material. Four of the product bands are fluorescent glycosides which have high Rf's in benzene/2-propanol (19:1). The major product is purified to homogeneity by two separate preparative layer chromatography procedures employing multiple developments in the above-named solvent system. The product (24 mg.) is a fluffy white solid after lyophilization from benzene. Mass spectral and 300 MHz nuclear magnetic resonance data are consistent with the structure of 4'-Q-(3,4-di-Q-acetyl-2,6-dideoxy-L-arabino-hexopyranosyl)-C-076 A_{2a} monosaccharide.

-30-

16024

EXAMPLE 10

4'-O-(2,4-di-O-acetyl-2,6-dideoxy-L-arabino-
hexopyranosyl C-076 B_{1a} monosaccharide-5-one

A 100 ml. round-bottomed flask is charged
5 with C-076 B_{1a} monosaccharide-5-one (30 mg.), 3,4-di-
O-acetyl-2,6-dideoxy-L-arabino-hexopyranosyl chloride
(diacetyl-oliviosyl chloride, 30 mg.), freshly
prepared silver oxide (400 mg.) and dry ether
(25 ml.). The stoppered reaction vessel is stirred
10 in the dark under nitrogen at ambient temperature
for 7 days when, by thin layer chromatography
[benzene/2-propanol (19:1)], the reaction appears
to make no further progress. Four additional 25 mg.
quantities of 3,4-di-O-acetyl-2,6-dideoxy-L-
15 arabino-hexopyranosyl chloride are added at 24 hour
intervals. During the reaction, the solids are
filtered, washed with ether, and discarded. The
filtrate is evaporated in vacuo and the colorless
gum is resolved by preparative layer chromatography
20 (above system, multiple development) into two major
bands: starting material and a slightly faster band,
which is chromatographically homogeneous and by
spectral analysis consistent with the 4'-O-(2,4-di-
O-acetyl-2,6-dideoxy-L-arabino-hexopyranosyl C-076
25 B_{1a} monosaccharide-5-one.

EXAMPLE 11

4"-O-benzoyl C-076 B_{1a} (from C-076 B_{1a} monosaccharide)
C-076 B_{1a} monosaccharide (35 mg.) is dissolved in dry ether and active manganese
5 dioxide (100 mg.) is added. The mixture is stirred magnetically for 18 hours at room temperature. The reaction mixture is filtered and the solids are washed with ether. The filtrate contains only one spot by thin layer chromatography [tetrahydro-
10 furan/chloroform (9:1)] and the solvent is evaporated in vacuo to afford B_{1a} monosaccharide-5-one as a colorless glass. Ether (25 ml.), 1,5-anhydro-4-O-benzoyl-2,6-dideoxy-3-O-methyl-L-arabino-hex-1-enitol (10 mg.) and pyridinium p-toluene-
15 sulfonate (1 mg.) are added and the stoppered vessel is stirred at ambient temperature for an additional 24 hour period. The volatiles are removed in vacuo and methanol (10 ml.) and sodium borohydride (20 mg.) are added to the residual
20 gum. A yellow color developed rapidly which faded within 15 minutes. After stirring for 2 hours under nitrogen, 3ml. of acetone are added and stirring is continued for an additional hour. The solvents are evaporated in vacuo and the residue
25 resolved on preparative layer chromatography [benzene/2-propanol (19:1)] to furnish the desired compound as the major product. Mass spectral and 300 Mhz nuclear magnetic resonance data are consistent with the structure of 4"-O-benzoyl
30 C-076 B_{1a}.

EXAMPLE 12

4'-Q- \underline{D} -glucopyranosyl C-076 A_{2a} monosaccharide
(Deprotection using methanolic ammonia)

5 4'-Q-(2,3,4,6-tetra-Q-acetyl- \underline{D} -gluco-
pyranosyl) C-076 A_{2a} monosaccharide (10 mg.) is
covered with 10 ml. of methanol which had previously
been saturated with anhydrous ammonia at 0°C. The
flask is stoppered tightly and stored at ambient
temperature for three days. Thin layer chromatography
10 monitoring [trichloromethane/methanol (9:1)] shows
no remaining peracetylated materials. The solvent
is evaporated under a stream of nitrogen and the
product is purified from minor contaminants by
preparative layer chromatography on silica gel
15 with the above solvent system as the eluent. The
homogeneous product is then lyophilized from
benzene affording pure 4'-Q-(β - \underline{D} -glucopyranosyl)C-076
A_{2a} monosaccharide.

EXAMPLE 13

20 5-Q-glucopyranosyl C-076 B_{1a}

(Deprotection using catalytic methoxide)

5-Q-(2,3,4,6-tetra-Q-acetyl- \underline{D} -gluco-
pyranosyl) C-076 B_{1a} (5 mg.) is dissolved in
anhydrous methanol (10 ml.) in a small flask that
25 has been flamed and cooled under nitrogen. A
small amount of sodium methoxide powder (3-5 mg.)
is added and the stoppered flask is stored at
ambient temperature overnight. A chunk of dry ice
is added to the reaction mixture before it was
30 evaporated under a stream of nitrogen. The product
is separated from minor impurities by preparative
layer chromatography on silica gel using di-
chloromethane/methanol (9:1) as the eluent.

5-O-glucopyranosyl C-076 B_{1a} is obtained in excellent yield as a lyophilized foam (from benzene).

EXAMPLE 14

5 Following the Koenigs-Knorr Method of Examples 3 or 8 using the appropriate starting materials, the following compounds are prepared:

- 5-O-peracetyl-L-olivosyl C-076 B_{1a}
- 5-O-peracetyl-L-olivosyl C-076 B_{1a} monosaccharide
- 10 5-O-peracetyl-L-rhamnopyranosyl C-076 B_{1a}
- 23-O-peracetyl-L-rhamnopyranosyl C-076 B_{2a}
- 5-O-peracetyl-L-rhamnopyranosyl C-076 B_{2a}
- 5-O-peracetyl-L-arabinopyranosyl C-076 B_{1a} mono-saccharide
- 15 23-O-peracetyl-L-arabinopyranosyl C-076 B_{2a} mono-saccharide
- 5-O-peracetyl-L-arabinopyranosyl C-076 B_{2a} mono-saccharide
- 5-O-peracetyl-L-lyxopyranosyl C-076 B_{1a}
- 20 23-O-peracetyl-L-lyxopyranosyl C-076 A_{2a} mono-saccharide
- 23-O-peracetyl-L-olivosyl C-076 A_{2a}
- 5-O-peracetyl-D-glucopyranosyl-22,23-dihydro C-076 B_{1a}
- 25 5-O-peracetyl-D-galactopyranosyl-22,23-dihydro C-076 B_{1a}
- 23-O-peracetyl-D-arabinopyranosyl C-076 A_{2a}
- 23-O-peracetyl-D-glucosaminyl C-076 A_{2a}
- 23-O-peracetyl-D-glucuronyl C-076 A_{2a}
- 30 5-O-peracetyl-D-glucuronyl C-076 B_{1a}
- 23-O-peracetyl-D-mannopyranosyl C-076 A_{2a}

EXAMPLE 15

Following the Koenigs-Knorr Method of Example 10, using the appropriate protected starting materials, and reducing the product 5 therefrom by the borohydride reduction method of Example 11, the following products are prepared:

- 13-O-peracetyl-L-olivosyl C-076 B_{1a} aglycone
- 13-O-peracetyl-D-maltosyl C-076 A_{1a} aglycone
- 13-O-peracetyl-D-glucosaminy C-076 B_{1a} aglycone
- 10 13-O-peracetyl-D-glucuronyl-22,23-dihydro C-076
B_{1a} aglycone
- 4'-O-peracetyl-L-lyxopyranosyl-22,23-dihydro C-076
B_{1a} monosaccharide

EXAMPLE 16

- 15 Following the orthoester method of Examples 1,2 or 7, using the appropriate starting materials, the following products are obtained.

- 4"-O-peracetyl-L-rhamnopyranosyl C-076 A_{2a}
- 4"-O-peracetyl-L-rhamnopyranosyl C-076 A_{1a}
- 20 4"-O-peracetyl-D-galactopyranosyl C-076 A_{2a}
- 4"-O-peracetyl-D-glucopyranosyl-22,23-dihydro C-076
B_{1a}
- 4'-O-peracetyl-L-lyxopyranosyl C-076 B_{1a} mono-
saccharide
- 25 4'-O-peracetyl-L-rhamnopyranosyl C-076 B_{1a} mono-
saccharide
- 4'-O-peracetyl-L-rhamnopyranosyl-22,23-dihydro
C-076 B_{1a} monosaccharide

EXAMPLE 17

Following the Helferich modification of Example 9, using the appropriate starting materials, the following products are obtained:

- 5 4'-Q-peracetyl-L-rhamnopyranosyl C-076 A_{2a} mono-saccharide
- 4'-Q-peracetyl-L-oliviosyl C-076 A_{2a} monosaccharide
- 13-Q-peracetyl-L-lyxopyranosyl C-076 A_{2a} aglycone
- 23-Q-peracetyl-L-lyxopyranosyl C-076 A_{2a} aglycone
- 10 13-Q-peracetyl-L-oliviosyl C-076 A_{2a} aglycone
- 23-Q-peracetyl-L-oliviosyl C-076 A_{2a} aglycone
- 13-Q-peracetyl-L-arabinopyranosyl C-076 A_{2a} aglycone
- 4'-Q-peracetyl-L-oleandrosyl-22,23-dihydro C-076 A_{1a} monosaccharide

15

EXAMPLE 18

Following the glycal method of Example 4, using the appropriate protected starting materials, and reducing the product therefrom by the borohydride method of Example 11, the following 20 products are obtained:

- 4'-Q(peracetyl-2-deoxy-D-arabino-hexopyranosyl) C-076 B_{1a} monosaccharide and 4'-Q-(peracetyl-2,3-dideoxy-D-erythro-hex-2-enopyranosyl) C-076 B_{1a} monosaccharide-from D-glucal
- 25 4'-Q-(peracetyl-2-deoxy-L-threo-pentopyranosyl) C-076 B_{1a} monosaccharide and 4'-Q-(peracetyl-23-dideoxy-L-glycero-pent-2-enopyranosyl) C-076 B_{1a} monosaccharide-from L-arabinal

-36-

16024

4'-Q-(peracetyl-2-deoxy-L-erythro-pentopyranosyl)
C-076 B_{1a} monosaccharide and 4'-Q-(peracetyl-
23-dideoxy-L-glycero-pent-2-enopyranosyl)
C-076 B_{1a} monosaccharide from L-xylal

5 13-Q-(peracetyl-2-deoxy-D-lyxo-hexopyranosyl)
C-076 B₁ aglycone and 13-Q-(peracetyl-2,3-
dideoxy-D-threo-hex-2-enopyranosyl) C-076
B_{1a} aglycone from D-galactal

10 4"-Q-isovaleryl-22,23-dihydro C-076 B_{1a} and 4"-Q-
isovaleryl-3"-deoxy-2,"3"-didehydro C-076
B_{1a} - from 4-Q-valeryl-L-oleandral

EXAMPLE 19CHROMATOGRAPHIC MOBILITIES OF PROTECTED C-076MACROLIDE GLYCOSIDES RELATED TO A_{2a}

ON 5 X 20 CM. SILICA GEL THIN LAYER

CHROMATOGRAPHY PLATES (DOUBLE DEVELOPMENT)BENZENE/2-PROPANOL (19:1)

<u>COMPOUND NAME</u>	
C-076 A _{2a} (reference)	0.23
23-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 A _{2a}	0.24
23-O-(2,2',3',4',6,6'-hepta-O-acetyl-maltosyl) C-076 A _{2a}	0.30
4"-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 A _{2a}	0.41
23-O-acetyl-4"-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 A _{2a}	0.40
A _{2a} monosaccharide (reference)	0.34
4'-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 A _{2a} monosaccharide	0.45
23-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 A _{2a} monosaccharide	0.33
4'-O-(3,4-di-O-acetyl-2,6-dideoxy-L-arabino-hexopyranosyl) C-076 A _{2a} monosaccharide (oliviosyl)	0.44
C-076 A _{2a} aglycone (reference)	0.38
23-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 A _{2a} aglycone	0.37
13-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 A _{2a} aglycone	0.26
13-O-(4-D-acetyl-2,3,6-trideoxy-α-L-erythro-Hex-2-enopyranosyl) C-076 A _{2a} aglycone	0.48
13,23-Bis-O-(4-O-acetyl-2,3,6-trideoxy-α-L-erythro-hex-2-eno- pyranosyl) C-076 A _{2a} aglycone	0.55

EXAMPLE 20

CHROMATOGRAPHIC MOBILITIES OF DEPROTECTED
C-076 MACROLIDE GLYCOSIDES
5 X 20 CM. SILICA GEL THIN LAYER CHROMATOGRAPHY
PLATES (SINGLE DEVELOPMENT)

COMPOUND NAME	DICHLOROMETHANE/METHANOL(9:1)
C-076 A ₂ (reference)	0.58
4"-O-D-glucopyranosyl C-076 A ₂	0.55
4'-O-D-glucopyranosyl C-076 A _{2a} monosaccharide	0.57
C-076 B ₂ (reference)	0.46
23-O-maltosyl C-076 B _{2a}	0.42
23-O-D-glucopyranosyl C-076 B _{2a} monosaccharide	0.47

EXAMPLE 21

CHROMATOGRAPHIC MOBILITIES OF PROTECTED
C-076 MACROLIDE GLYCOSIDES
RELATED TO B_{1a} OR B_{2a} 5 X 20 SILICA GEL THIN
LAYER CHROMATOGRAPHY PLATES (DOUBLE DEVELOPMENT)

BENZENE/2-PROPANOL (19:1)

COMPOUND NAME

C-076 B _{1a} (reference)	0.18
5-O-acetyl-4"-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 B _{1a}	0.33
5-O-acetyl-4"-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 B _{1a}	0.32
5-O-(2,3,4,6-tetra-O-acetyl-D-galactopyranosyl) C-076 B _{1a}	0.26
4'-O-(3,4-di-O-acetyl-2,6-dideoxy-L-arabino hexopyranosyl)- C-076 B _{1a} monosaccharide-5-ona (olivosyl)	0.41
5-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 B _{1a} monosaccharide	0.21
C-076 B _{2a} (reference)	0.15
23-O-(2,2',3,3',4',6,6'-hepta-O-acetyl-maltosyl) C-076 B _{2a}	0.18
23-O-(2,3,4,6-tetra-O-acetyl-D-glucopyranosyl) C-076 B _{2a}	0.20

PREPARATION 1Glycal synthesis
4-O-benzoyl-L-oleandral

A mixture of 40% sodium hydroxide (75 ml.)
5 diethyl ether (200 ml.) are stirred in an ice bath,
while 16.0 g. of nitrosomethylurea is added in portions
over a period of 20 min. The mixture is stirred for
an additional hour in the ice bath and the ether
solution then decanted and dried over sodium hydroxide
10 pellets.

To a solution of L-rhamnal (Methods in Car-
bohydrate Chemistry, Vol. II, p. 407-8), 2.1 g. in
diethyl ether (200 ml.), is added stannous chloride
dihydrate (200 mg.) in 20 ml. of ether followed by about
15 one third of the diazomethane solution. The reaction
mixture remains at room temperature (23°C.) and after
30 min. another 200 mg. of the diazomethane solution.
The reaction is stirred overnight (18 hrs.) and
silica gel t/c (dichloromethane/methanol, 19:1) shows
20 the reaction to be essentially complete. Acetic acid
is added to dispose the yellow color and the mixture
is filtered through Supercel. The filtrate is washed
with 10% sodium bicarbonate solution and then with
saturated salt solution. The solution is dried and
25 evaporated to a residue, which is then purified to the
homogeneity by silica gel chromatography (dichloromethane/
rhamnal) as a colorless syrup (77%).

-41-

16024

The methylation product (2.23 g) is dissolved in pyridine (20 ml.) and stirred with cooling while 2.25 ml. of benzoyl chloride is added. The temperature rises slightly and a light precipitate forms. The cooling bath is removed after two hours and stirring maintained for 48 hrs. Water is then added (15 drops) and after 45 min., 200 ml. of dichloromethane. The mixture is washed with 1N HCl until the extracts are acidic and then with sodium bicarbonate solution and finally saturated salt solution. The organic phase is dried and the solvent removed in vacuo to furnish a colorless syrup, which could be purified further by silica gel chromatography using (99:1) dichloromethane/tetrahydrofuran as the eluant to furnish 3.6 g of protected glycal.

The product contains a small amount of the 4-0-methyl isomer, which could be resolved by silica gel high performance liquid chromatography using 7.5% ethyl acetate in n-hexane as eluant.

20

PREPARATION 2

Of glycosylhalide synthesis

L-oliviosyl chloride(2,6-dideoxy-3,4-di-0-acetyl-L-arabino-hexopyranosyl chloride

25 Dry hydrogen chloride gas is bubbled slowly into a benzene solution of 3,4-di-0-acetyl-L-rhamnal (7.5 g. in 50 ml. benzene). The solution is cooled in an ice salt bath. When the solution is saturated (about 20 min.), the addition of hydrogen chloride is discontinued and the mixture allowed to stand at 0°C. 30 for one hour. The solvent is then removed under vacuum and toluene (50 ml.) added and removed in vacuo

(high vacuum) three times. The residual oil is dissolved in ether at 0°C and petroleum ether (30-60°C) added to induce crystallization. The colorless crystals are filtered under nitrogen
5 washed with petroleum ether and dried on the funnel under a stream of nitrogen before storage at -5°C (6.8 g.).

PREPARATION 3

C-076 Ala Aglycone

10 100 Mg. of C-076 Ala is dissolved in 5 ml. of dioxane, stirred and added at room temperature to a mixture of 0.1 ml. of concentrated sulfuric acid, 1.9 ml. of methanol and 3.0 ml. of dioxane. The reaction mixture is stirred overnight at room
15 temperature. 473 Mg. of solid sodium bicarbonate is added and the mixture stirred for 20 minutes. 3 ml. of water is added and stirred for an additional 10 minutes. The reaction mixture is concentrated and 40 ml. of chloroform is added and shaken. The aqueous
20 layer is separated and extracted with 5 ml. of chloroform. The organic layers are combined and washed once with dilute sodium chloride solution, dried over magnesium sulfate and evaporated to dryness in vacuo. 1/2 of the residue is placed on 5 preparative
25 layer chromatography silica gel plates and eluted with 2% methanol in chloroform affording 4 bands of material. The remainder of the material is run on 2 preparative layer chromatography plates eluting with 2% methanol in chloroform affording 4 band similar to
30 the first series. The second fastest bands are removed from each of the plates combined, extracted and evaporated to dryness in vacuo, and rechromatographed on a preparative layer chromatography silica gel plate eluting with 3% tetrahydrofuran and chloroform
35 affording 9.4 mg. of a fluffy white solid which is identified by mass spectrometry as C-076 Ala aglycone.

PREPARATION 4C-076 A2a Aglycone

2 G. of C-076 A2a is combined with 40 ml. of
a 1:1 (volume/volume) solution of concentrated sulfuric
5 acid in methanol. The reaction mixture is stirred at
room temperature for 17 hours and diluted with 300 ml.
of chloroform. The mixture is washed once with 30 ml.
of saturated sodium bicarbonate solution, once with
30 ml. saturated sodium chloride solution, dried over
10 magnesium sulfate and evaporated to dryness in vacuo.
5 ml. of methanol is added to the residue and allowed
to stand at room temperature overnight. Cooling of
the mixture in ice causes the slow precipitation of
crystals. A supernatant is removed and the solid
15 crystals washed twice with 1 ml. of cold methanol
affording 340 mg. of a white solid. The mother liquor
and washings are evaporated down to a volume of about
2 ml. and allowed to stand affording an additional
crop to crystals. 630 mg. of a white solid is obtained
20 which is combined with the first batch of crystals
and 8 ml. of methanol and evaporated to a volume of
2.5 ml. and allowed to stand for several hours.
910 mg. of an off white solid is obtained which mass
spectrometry identifies as C-076 A2a aglycone.

25

PREPARATION 5C-076 A2a Monosaccharide

500 mg. of C-076 A2a is dissolved in 10 ml.
of a solution of 0.1 ml. of concentrated sulfuric acid
and 9.9 ml. of isopropanol. The reaction mixture is
30 stirred at room temperature overnight. 125 ml. of
chloroform is added and the mixture washed once with
10 ml. of saturated sodium bicarbonate and once with
10 ml. of water. The organic layer is dried over
magnesium sulfate and evaporated to dryness in vacuo

affording a pale yellow solid material which is dissolved in chloroform and placed on 5 preparative layer chromatography silica gel plates and eluted twice with 2% benzene in ethylacetate. The slower moving major fraction contains 367 mg. of a white powder after lyophilization from benzene which mass spectrometry and 300 MHz nuclear magnetic resonance indicates is C-076 A2a monosaccharide.

PREPARATION 6

10 C-076 Bla Monosaccharide and C-076 Bla Aglycone

2.5 Ml. of a solution consisting of 0.5 ml. of water 0.5 ml. concentrated sulfuric acid and 9.0 ml. of dioxane is added and the reaction mixture stirred at room temperature for 17 hours. 50 Ml. of ether is added followed by 25 ml. of saturated aqueous sodium bicarbonate solution. The two layer mixture is shaken, the aqueous layer separated and the organic layer washed with water, dried and evaporated to dryness in vacuo. Benzene is added to the residue and the benzene layer is dried and lyophilized affording 60 mg. of yellow material. The material is placed on a preparative layer chromatography silica gel plate and eluted with chloroform-tetrahydrofuran in the volume ratio of 9:1 and 2 bands are observed with an Rf of 0.15 and 0.35. 300 MHz nuclear magnetic resonance identifies the two spots as C-076 Bla monosaccharide and C-076 Bla aglycone respectively. 16 Mg. of each fraction is obtained.

PREPARATION 7

30 C-076 Bla Monosaccharide

100 Mg. of C-076 Bla is dissolved in 5.0 ml. of tetrahydrofuran and stirred at room temperature while 5.0 ml. of a cold aqueous solution of 10% sulfuric

-45-

16024

acid (volume/volume) is added dropwise with stirring. The reaction mixture is stirred at room temperature for 18 hours. 75 ml. of methylene chloride and 25 ml. of saturated aqueous sodium bicarbonate is added and the layers shaken and separated. The organic layer is washed with aqueous sodium chloride solution and an equal volume of water. The organic layer is dried and evaporated to dryness in vacuo affording 70 mg. of a colorless oil. High pressure liquid identifies the residual oil as C-076 B1a monosaccharide.

PREPARATION 8

C-076 B2a Aglycone

2 G. of C-076 B2a is combined with 40 ml. of a 1% solution of concentrated sulfuric acid in methanol (volume/volume). The reaction mixture is stirred at room temperature for 17 hours. 300 ml. of chloroform is added followed by 30 ml. of an aqueous saturated sodium bicarbonate solution. The layers are separated and the organic layer washed with 30 ml. of saturated sodium chloride solution, dried over magnesium sulfate and evaporated to dryness in vacuo. 5 ml. of methanol is added to dissolve the residue and the mixture allowed to stand at room temperature and then cooled in an ice bath, whereupon crystallization occurs. The supernatant is removed and the residue washed twice with 1 ml. portions of cold methanol and the solid crystals dried overnight and then in vacuo at 35°C affording 1.0 g. of white crystals. A second crop is obtained by evaporating the mother liquors to a volume of 2 ml. and allowing to stand overnight at room temperature. 2 ml. of methanol is added and the mixture aged in an ice bath affording 140 mg. of a yellow solid. The two solid fractions are combined and dissolved in boiling methanol, about

30 ml. of methanol is required. The solution is filtered hot and concentrated to a volume of about 20 ml. in vacuo whereupon solids begin to precipitate. The solution is filtered hot and the solid materials washed with methanol affording 340 mg. of a white solid. The filtrates are boiled down to a volume of about 8 ml. and set aside to crystallize at room temperature affording 433 mg. of a white solid. Mass spectrometry shows the two fractions to be identical and to be identified as C-076 B2a aglycone.

PREPARATION 9

C-076 B2a Monosaccharide and C-076 B2a Aglycone

20 Mg. of C-076 B2a is combined with 4 ml. of a solution prepared by combining 0.1 ml. of concentrated sulfuric acid and 9.9 ml. of isopropanol. The reaction mixture is stirred at room temperature for 16 hours, 189 mg. of sodium bicarbonate is added followed by a few drops of water. The volume is reduced to about 1/2 and 30 ml. of chloroform and 3 ml. of water is added and the mixture shaken. The layers are separated and the aqueous layer extracted with an additional 5 ml. of chloroform. The organic layers are combined, washed once with dilute sodium chloride solution, dried over sodium sulfate and magnesium sulfate and evaporated to dryness in vacuo. The residue is placed on two preparative layer silica gel chromatography plates and eluted twice with 5% tetrahydrofuran in chloroform. 4 Bands of material are observed and individually removed from the preparative chromatography plates. The slowest band affords 7.3 mg. of a white solid which is identified by mass spectrometry as C-076 B2a monosaccharide. The next slowest band affords 1.3 mg. of a white solid and it is identified by mass spectrometry as C-076 B2a aglycone.

PREPARATION 1022,23-Dihydro C-076 Ala

5 51.0 Mg. of C-076 Ala and 14.4 mg. of tris
(triphenylphosphine)rhodium (I) chloride are combined
in 3.5 ml. of benzene and hydrogenated for 20 hours at
room temperature under atmospheric pressure. The crude
reaction mixture is chromatographed on a preparative
layer chromatography plate eluting twice with 10%
tetrahydrofuran in chloroform. The product is removed
10 from the support using ethyl acetate which is
evaporated to dryness and the residue analyzed with
300 MHz nuclear magnetic resonance and mass
spectroscopy indicating the preparation of 22,23-
dihydro C-076 Ala.

PREPARATION 1122,23-Dihydro C-076 Bla

15 A solution of 1.007 g. of C-076 Bla, 314 mg. of
tris(triphenylphosphine)rhodium(I)chloride and 33 ml.
of benzene is hydrogenated for 21 hours at room
20 temperature under 1 atmosphere of hydrogen pressure.
The solvent is removed in vacuo and the residue
dissolved in a 1:1 mixture of methylene chloride and
ethyl acetate and filtered. The filtrate is placed
on a column of 60 g. of silica gel eluting with a 1:1
25 mixture of methylene chloride and ethyl acetate taking
10 ml. fractions. Fractions 14-65 are combined and
evaporated to dryness affording 1.118 g. of a solid
material which is indicated by high pressure
liquid chromatography to be a 60/40 mixture of the
30 hydrogenated product and starting material. The mixture
is rehydrogenated in 55 ml. of benzene adding 310 mg. of
tris (triphenylphosphine) rhodium (I) chloride and
stirring for 21 hours at room temperature under 1

atmosphere of hydrogen pressure. The solvent is removed in vacuo and the residue chromatographed on 80 g. of silica gel using 40:60 mixture of ethyl acetate and methylene chloride as eluant. 10 Ml. 5 fractions are taken and the product appears in fractions 26-80. These fractions are combined and evaporated to dryness in vacuo affording a yellow oil. The oil is dissolved in benzene and lyophilized affording a pale yellow powder which is identified 10 as 22,23-dihydro C-076 Bla by mass spectrometry and 300 MHz nuclear magnetic resonance. 0.976 G. of product is obtained.

PREPARATION 12

22,23-Dihydro C-076 Bla Monosaccharide

15 395 Mg. of 22,23-dihydro C-076 Bla is added to a stirred solution of 50 ml. of 1% sulfuric acid in isopropanol and the solution is stirred for 14 hours at room temperature. The reaction mixture is treated as in Example 4 affording 0.404 g. of a foam after 20 lyophilization from benzene. The foam is chromatographed on 6 preparative layer silica gel chromatography plates eluting twice with 4% tetrahydrofuran in chloroform. The monosaccharide with a R_f 0.15 is collected and washed from the silica gel with a total 25 of 650 ml. of ethyl acetate. The combined washings are evaporated to dryness and the residue lyophilized from benzene to afford 0.2038 g. of 22,23-dihydro C-076 Bla monosaccharide which high pressure liquid chromatography indicates to be essentially pure.

PREPARATION 1322,23-Dihydro C-076 Bla Aglycone

0.486 G. of 22,23-dihydro C-076 Bla is added to a stirred solution of 50 ml. of 1% sulfuric acid in methanol and the reaction mixture stirred for 13 hours at room temperature. The reaction mixture is diluted with 250 ml. of methylene chloride and washed with 50 ml. of saturated aqueous potassium bicarbonate and 50 ml. of water. The aqueous layer is washed twice with 20 ml. portions of methylene chloride and the combined organic phases are dried with saturated brine and sodium sulfate and evaporated to dryness in vacuo affording 0.480 g. of a pale yellow foam. The foam is dissolved in 4 ml. of methylene chloride and placed on 4 preparative layer chromatography silica gel plates and eluted 4 times with 4% tetrahydrofuran and chloroform. The product is recovered from the silica gel plates affording an oily residue which is lyophilized from benzene affording 255.8 mg. of a white solid. Traces of methyl oleandroside are indicated to be present in the solid material. The white solid is then lyophilized again from benzene and placed under high vacuum for 20 hours to remove the impurity affording 22,23-dihydro C-076 Bla aglycone.

PREPARATION 14A. C-076 Ala 4"-O-Acetate

A solution of 27 mg. of 4-dimethylamino-pyridine in 1 ml. of methylene chloride is prepared and a separate solution of 0.208 ml. of acetic anhydride in 10 ml. of methylene chloride is prepared. 0.5 ml. of

each solution is added to 10 mg. of C-076 Ala, mixed well and allowed to stand at room temperature overnight. The reaction mixture is diluted to 4 ml. with methylene chloride and 0.5 ml. of water is added and shaken. The layers are separated and the organic layer is dried over magnesium sulfate and evaporated to dryness under a stream of nitrogen. Benzene is added and the solution is lyophilized affording 10 mg. of an off-white fluffy solid.

10 Preparative layer chromatography on silica gel eluting with 10% tetrahydrofuran in chloroform affords 8.2 mg. of a fluffy, off-white solid, which nuclear magnetic resonance and mass spectrographic analysis reveals to be C-076 Ala 4"-O-acetate.

15 B. C-076 A2a 4"-O-Acetate

Following the above procedure 5 mg. of C-076 A2a is acetylated affording 4.4 mg. of a product which is demonstrated by mass spectrometry and nuclear magnetic resonance to be C-076 A2a

20 4"-O-acetate.

C. C-076 A2a 4",23-di-O-acetate

10 Mg. of C-076 A2a is acetylated in 0.5 ml. of pyridine and 0.25 ml. acetic anhydride at 100°C for 2 hours. The reaction mixture is worked up

25 using preparative layer chromatography on silica gel as previously described affording 5.9 mg. of a fluffy white solid which mass spectrometry and nuclear magnetic resonance reveal to be C-076 A2a 4",23-O-acetate.

PREPARATION 15C-076 A2a 4" O-Propionate

25 Mg. of C-076 A2a is combined with 15 drops of dry pyridine and cooled in ice while 5 drops of propionic anhydride is added. The reaction mixture is stoppered, mixed well and allowed to stand in an ice bath overnight. The reaction mixture is diluted with ether and benzene and shaken with some ice water. The layers are separated and the organic layer is dried over magnesium sulfate. The solvent is evaporated under a stream of nitrogen, benzene is added and the solution is lyophilized affording 20 mg. of a white solid. Preparative layer chromatography on silica gel eluting with 5% tetrahydrofuran in chloroform affords 16.6 mg. of a white solid which is analysed by nuclear magnetic resonance and mass spectrometry as C-076 A2a 4"-O-propionate.

PREPARATION 16C-076 Bla 4",5-di-O-acetate


20 Following the procedure of Preparation 14, 5.2 mg. of C-076 Bla is acetylated with 10 drops of pyridine and 6 drops of acetic anhydride affording, after preparative layer chromatography on silica gel and lyophilization, 5.2 mg. of a white fluffy solid which mass spectrometry indicates is C-076 Bla 4",5-di-O-acetate.

PREPARATION 17C-076 Bla 4",O-Acetate and C-076 Bla 4",5-Di-O-Acetate


20 Mg. of C-076 Bla is dissolved in 12
drops of pyridine, cooled in an ice bath and combined
5 with 4 drops of acetic anhydride. The reaction
mixture is maintained in an ice bath for 2 1/2 hours,
chilled benzene is added, the reaction mixture
freeze dried and the solid material chromatographed
on silica gel plates eluting with 10% isopropanol in
10 benzene. The product with the highest Rf is
identified by mass spectrometry as C-076 Bla 4",
5-di-O-acetate, 4.7 mg. is obtained. The next most
advanced spot is identified by mass spectrometry as
C-076 Bla 4",O-acetate; 9.3 mg. is obtained.

CLAIMS

1. A compound having the formula :



in which the broken line indicates that a single or double bond is present in the 22(23) position; R_1 is hydroxy, loweralkanoyloxy, or glycosyloxy, and is present only when the broken line indicates a single bond; R_2 is iso-propyl or sec-butyl; R_3 is hydrogen, methyl, loweralkanoyl or glycosyl; and R is hydrogen, lower alkanoyl, glycosyl,




where R_4 is hydrogen, loweralkanoyl or glycosyl; in which the glycosyl groups are polyhydroxy (5 or 6)-membered cyclic acetals that are optionally substituted with lower alkyl and in which the hydroxy groups may be optionally substituted with lower alkyl or lower alkanoyl and their mono-unsaturated derivatives; provided that at least one of R , R_1 , R_3 or R_4 must contain a glycosyl group, and that if only R contains a glycosyl group, that group is other than α -L-oleandrosyl or α -L-oleandrosyl- α -L-oleandrosyl.


2. A compound as claimed in Claim 1, in which the glycosyl residue of R , R_1 , R_3 , or R_4 is a monosaccharide, disaccharide or trisaccharide in which the saccharide residue(s), which may be the same or different in the disaccharides and trisaccharides, is/are glucopyranosyl, galactopyranosyl, mannopyranosyl, maltosyl, arabinopyranosyl, lyxopyranosyl, xylopyranosyl, ribopyranosyl, oleandrosyl, rhamnopyranosyl, fucopyranosyl, lactosyl, ribofuranosyl, mannofuranosyl, glucofuranosyl, arabinofuranosyl, mycarosyl, cladinosyl, desosaminosyl, daunosaminosyl, mycaminosyl, cymarosyl and/or olivosyl.

3. A compound as claimed in Claim 2, in which the glycosyl residue is a monosaccharide or disaccharide in which the saccharide residue(s), which may be the same or different in the disaccharide, is/are glucopyranosyl, rhamnopyranosyl, oleandrosyl, and/or daunosaminosyl.

4. A process for the preparation of a compound as claimed in Claim 1 comprising reacting a compound having the formula :



in which the broken line indicates that a single or double bond is present in the 22(23) position; R_1 is hydroxy, loweralkanoyloxy, or glycosyloxy, and is present only when the broken line indicates a single bond; R_2 is iso-propyl or sec-butyl; R_3 is hydrogen, methyl, loweralkanoyl or glycosyl; and R' is hydrogen, lower-alkanoyl, glycosyl,



where R'_4 is hydrogen, loweralkanoyl or glycosyl; in which the glycosyl groups are polyhydroxy (5 or 6)-membered cyclic acetals that are optionally substituted with lower alkyl and in which the hydroxy groups may be optionally substituted with lower alkyl or lower alkanoyl and their mono-unsaturated derivatives; provided that at least one of R , R_1 , R_3 and R_4 must contain a free hydroxy

group, with a protected halohexose or protected halopentose in the presence of silver oxide.

5. A process for the preparation of a compound as claimed in Claim 1 comprising reacting a compound having the formula set forth in Claim 4 with a protected halohexose or protected halopentose in the presence of mercuric oxide or mercuric cyanide.

6. A process for the preparation of a compound as claimed in Claim 1 comprising reacting a compound having the formula set forth in Claim 4 with a protected halohexose or protected halopentose in the presence of silver trifluoromethyl sulfonate.

7. A process for the preparation of a compound as claimed in Claim 1 comprising reacting a compound having the formula set forth in Claim 4 with the orthoester prepared from a loweralkanol and a protected halohexose or protected halopentose in the presence of a catalytic amount of mercuric bromide or mercuric chloride.

8. A process for the preparation of a compound of Claim 1, in which the glycosyl group is a hexose that is unsubstituted at the 3-position or has a 2,3-unsaturation, comprising reacting a compound having the formula set forth in Claim 4 with a 1,2-unsaturated hexose, in the presence of a catalytic amount of a Lewis acid.

9. A composition useful for the treatment of parasitic infections comprising a compound as claimed in Claim 1 and an inert carrier.

10. The use of a compound as claimed in Claim 1 in the treatment of parasitic infections.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

Application number

EP 79 30 1514

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
	<p>THE JOURNAL OF ANTIBIOTICS, vol. XXIX, 1976, no. 3, Japan Antibiotics Research Association, Tokyo Milbemycin $\beta_1, \beta_2, \beta_3$.</p> <p>* Pages 14,15,16 *</p> <p>--</p> <p>THE JOURNAL OF ANTIBIOTICS, Vol. XXIX, 1976, no. 6, Japan Antibiotics Research Association, Tokyo, Milbemycin $\alpha_1 - \alpha_{10}$.</p> <p>* Pages 35,36,37,38,39,40,41,42 *</p> <p>--</p> <p>US - A - 4 093 629 (M.H. FISHER)</p> <p>* Columns 1,2 *</p> <p>--</p> <p>FR - A - 2 178 778 (SANKYO CO.LTD.)</p> <p>* Pages 20-27 *</p> <p>--</p>	<p>1,9</p> <p>1,9</p> <p>1,9</p> <p>1,9</p>	<p>C 07 H 17/08 A 61 K 31/70// A 01 N 9/28</p> <p>TECHNICAL FIELDS SEARCHED (Int. Cl.)</p> <p>C 07 H 17/08 A 61 K 31/70</p>
INCOMPLETE SEARCH			CATEGORY OF CITED DOCUMENTS
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-9</p> <p>Claims searched incompletely: 10</p> <p>Claims not searched: 10</p> <p>Reason for the limitation of the search: Method for treatment of the human or animal body by surgery or therapy (See article 52(4) of the European Patent Convention)</p>			<p>X: particularly relevant</p> <p>A: technological background</p> <p>O: non-written disclosure</p> <p>P: intermediate document</p> <p>T: theory or principle underlying the invention</p> <p>E: conflicting application</p> <p>D: document cited in the application</p> <p>L: citation for other reasons</p> <p>&: member of the same patent family.</p> <p>corresponding document</p>
Place of search	Date of completion of the search	Examiner	
The Hague	04-10-1979	VERHULST	



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application number

EP 79 30 1514

-2-

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 1)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
	<p><u>FR - A - 2 348 970</u> (MERCK & CO.)</p> <p>* Pages 44-47 *</p> <p>--</p>	1,9	
P	<p><u>FR - A - 2 387 231</u> (MERCK & CO.)</p> <p>* Pages 18-21 *</p> <p>--</p>	1,9	
P	<p><u>EP - A - 1 688</u> (MERCK & CO.)</p> <p>* Pages 49-52 *</p> <p>--</p>	1,9	TECHNICAL FIELDS SEARCHED (Int. Cl. 1)
P	<p><u>EP - A - 1 689</u> (MERCK & CO.)</p> <p>* Pages 30-32 *</p> <p>----</p>	1,9	